

Practice Problems 13 : Ratio and Root tests, Leibniz test

1. Determine the values of  $\alpha \in \mathbb{R}$  for which  $\sum_{n=1}^{\infty} \left(\frac{\alpha n}{n+1}\right)^n$  converges.
2. Consider  $\sum_{n=1}^{\infty} a_n$  where  $a_n > 0$  for all  $n$ . Prove or disprove the following statements.
  - (a) If  $\frac{a_{n+1}}{a_n} < 1$  for all  $n$  then the series converges.
  - (b) If  $\frac{a_{n+1}}{a_n} > 1$  for all  $n$  then the series diverges.
3. Show that the series  $\frac{1}{1^2} + \frac{1}{2^3} + \frac{1}{3^2} + \frac{1}{4^3} + \frac{1}{5^2} + \frac{1}{6^3} + \dots$  converges and that the root test and ratio test are not applicable.
4. Consider the rearranged geometric series  $\frac{1}{2} + 1 + \frac{1}{8} + \frac{1}{4} + \frac{1}{32} + \frac{1}{16} + \frac{1}{128} + \frac{1}{64} \dots$ . Show that the series converges by the root test and that the ratio test is not applicable.
5. Consider the series  $\frac{1}{2} + \frac{1}{3} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{2^3} + \frac{1}{3^3} + \frac{1}{2^4} + \frac{1}{3^4} + \dots$ . Show that the ratio test is not applicable. Further show that  $(a_n)^{\frac{1}{n}}$  does not converge and that the root test given in Theorem 8 is applicable.
6.
  - (a) If  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  converges absolutely, show that  $\sum_{n=1}^{\infty} a_n b_n$  converges absolutely.
  - (b) If  $\sum_{n=1}^{\infty} a_n$  converges absolutely and  $(b_n)$  is a bounded sequence then  $\sum_{n=1}^{\infty} a_n b_n$  converges absolutely.
  - (c) Give an example of a convergent series  $\sum_{n=1}^{\infty} a_n$  and a bounded sequence  $(b_n)$  such that  $\sum_{n=1}^{\infty} a_n b_n$  diverges.
7. In each of the following cases, discuss the convergence/divergence of the series  $\sum_{n=1}^{\infty} a_n$  where  $a_n$  equals:
 

(a) $\frac{n!}{e^{n^2}}$	(b) $\frac{n^2 2^n}{(2n+1)!}$	(c) $(1 - \frac{1}{n})^{n^2}$	(d) $\frac{n^2}{3^n} (1 + \frac{1}{n})^{n^2}$
(e) $\sin\left(\frac{(-1)^n}{n^p}\right), p > 0$	(f) $(-1)^n \frac{(\ln n)^3}{n}$	(g) $(-1)^n \left(n^{\frac{1}{n}} - 1\right)^n$	(h) $\frac{2^n + n^2 - \ln n}{n!}$
(i) $\frac{\cos(\pi n) \ln n}{n}$	(j) $(1 + \frac{2}{n})^{n^2 - \sqrt{n}}$	(k) $\frac{n^2(2\pi + (-1)^n)^n}{10^n}$	
8. (\*) Let  $a_n \in \mathbb{R}$  and  $a_n > 0$  for all  $n$ .
  - (a) If  $\frac{a_{n+1}}{a_n} \leq \lambda$  eventually for some  $\lambda > 0$  then show that  $a_n^{\frac{1}{n}} \leq \lambda + \epsilon$  eventually for every  $\epsilon > 0$ . Observe that if the ratio test (Theorem 7) gives the convergence of a series then the root test (Theorem 8) also gives the convergence, but the converse is not true (why?).
  - (b) If  $\lim_{n \rightarrow \infty} a_n^{\frac{1}{n}} = \alpha$  and  $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = \beta$ , show that  $\alpha \leq \beta$ .
9.
  - (a) (\*) **(Dirichlet test)** Let  $\sum_{n=1}^{\infty} a_n$  be a series whose sequence of partial sums is bounded. Let  $(b_n)$  be a decreasing sequence which converges to 0. Show that  $\sum_{n=1}^{\infty} a_n b_n$  converges. Observe that Leibniz test is a particular case of the Dirichlet test.
  - (b) (\*) **(Abel's test)** Let  $\sum_{n=1}^{\infty} a_n$  be a convergent series and  $(b_n)$  be a monotonic convergent sequence. Show that  $\sum_{n=1}^{\infty} a_n b_n$  converges.
  - (c) Show that the series  $1 - \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{6} - \frac{1}{7} + \dots$  converges whereas the series  $1 + \frac{1}{2} - \frac{1}{3} + \frac{1}{4} + \frac{1}{5} - \frac{1}{6} + \frac{1}{7} + \frac{1}{8} - \dots$  diverges.
  - (d) Show that the series  $\sum_{n=1}^{\infty} \frac{(-1)^n}{n} (1 + \frac{1}{n})^n$ ,  $\sum_{n=1}^{\infty} \frac{(-1)^n}{n} \cos \frac{1}{n}$  and  $\sum_{n=1}^{\infty} (-1)^n \frac{\tan^{-1} n}{\sqrt{n}}$  converge.

Practice Problems 13 : Hints/Solutions

1. Since  $|\frac{an}{n+1}| \rightarrow \alpha$ , by the root test, the series converges for  $|\alpha| < 1$  and diverges for  $|\alpha| > 1$ . For  $|\alpha| = 1$ , the series diverges because  $(\frac{n}{n+1})^n \rightarrow \frac{1}{e} \neq 0$ .
2. (a) For  $a_n = \frac{1}{n}$ ,  $\frac{a_{n+1}}{a_n} < 1$  for all  $n$  but  $\sum_{n=1}^{\infty} \frac{1}{n}$  diverges.  
 (b) If  $\frac{a_{n+1}}{a_n} > 1$  then  $a_n \not\rightarrow 0$ . Hence  $\sum_{n=1}^{\infty} a_n$  diverges.
3. By the comparison test (with  $\frac{1}{n^2}$ ) the series converges.
4. The  $n$ th term  $a_n$  is  $\frac{1}{2^n}$  if  $n$  is odd and  $\frac{1}{2^{n-2}}$  if  $n$  is even. Since the consecutive ratio alternate in value between  $\frac{1}{8}$  and 2, the ratio test is not applicable. However  $a_n^{\frac{1}{n}} \rightarrow \frac{1}{2}$ .
5. Observe that  $(\frac{3}{2})^n$  and  $(\frac{2}{3})^n$  are subsequences of  $(\frac{a_{n+1}}{a_n})$ ; and  $(\frac{3}{2})^n \rightarrow \infty$   $(\frac{2}{3})^n \rightarrow 0$ . Therefore the ratio test (Theorem 7) is not applicable. But  $a_n$  can be either  $3^{-\frac{n}{2}}$  or  $2^{-\frac{n+1}{2}}$ . Since  $(3^{-\frac{n}{2}})^{\frac{1}{n}} \rightarrow \frac{1}{\sqrt{3}}$  and  $(2^{-\frac{n+1}{2}})^{\frac{1}{n}} \rightarrow \frac{1}{\sqrt{2}}$ ,  $a_n^{\frac{1}{n}} < L$  eventually for some  $L$  satisfying  $\frac{1}{\sqrt{2}} < L < 1$ . Hence the root test (Theorem 8) is applicable and the series converges.
6. (a) Since  $b_n \rightarrow 0$ ,  $|a_n b_n| \leq |a_n|$  eventually. Use the comparison test.  
 (b) Let  $|b_n| \leq M$  for some  $M$ . Then  $|a_n b_n| \leq M|a_n|$ . Use the comparison test.  
 (c) Consider  $a_n = \frac{(-1)^n}{n}$  and  $b_n = (-1)^n$ .
7. (a) Converges by the Ratio test.  
 (b) Converges by the Ratio test.  
 (c) Converges by the Root test:  $(1 - \frac{1}{n})^n \rightarrow \frac{1}{e}$  (If  $y = (1 - \frac{1}{n})^n$  then  $\ln y = \frac{\ln(1 - \frac{1}{n})}{\frac{1}{n}} \rightarrow -1$ ).  
 (d) Converges by the Root test:  $a_n^{\frac{1}{n}} \rightarrow \frac{e}{3} < 1$ .  
 (e) Converges by Leibniz test:  $\sin(\frac{(-1)^n}{n^p}) = (-1)^n \sin(\frac{1}{n^p})$ .  
 (f) Converges by Leibniz test: If  $f(x) = \frac{(\ln x)^3}{x}$  then  $f'(x) < 0$  for all  $x > e^3$ .  
 (g) Converges absolutely by the Root test.  
 (h) Converges: By the LCT test with  $\frac{2^n}{n!}$  and then the Ratio test for  $\sum_{n=1}^{\infty} \frac{2^n}{n!}$ .  
 (i) Converges by Leibniz test:  $\cos(\pi n) = (-1)^n$ .  
 (j) Diverges because  $(1 + \frac{2}{n})^{n^2 - \sqrt{n}} \not\rightarrow 0$  as  $(1 + \frac{2}{n}) > 1$ .  
 (k) Converges absolutely: Use  $|a_n| \leq \frac{n^2(2\pi+1)^n}{10^n}$  and then the Ratio test.
8. (a) Suppose  $\frac{a_{n+1}}{a_n} \leq \lambda$  for all  $n \geq N$  for some  $N$ . Then for all  $n \geq N$ ,

$$a_n = \frac{a_n}{a_{n-1}} \frac{a_{n-1}}{a_{n-2}} \dots \frac{a_{N+1}}{a_N} a_N \leq \lambda^{n-N} a_N.$$

Therefore  $a_n^{\frac{1}{n}} = (\lambda^{1 - \frac{N}{n}}) a_N^{\frac{1}{n}} \leq \lambda + \epsilon$  eventually for any  $\epsilon > 0$  as  $a_N^{\frac{1}{n}} \rightarrow 1$ .

Suppose the Ratio test (Theorem 7) implies the convergence of a series  $\sum_{n=1}^{\infty} a_n$ . Then there exists a  $\lambda$  such that  $0 < \lambda < 1$  and  $\frac{a_{n+1}}{a_n} \leq \lambda$  eventually. Then, by the previous part,  $a_n^{\frac{1}{n}} \leq \lambda + \frac{(1-\lambda)}{2} < 1$  eventually. Therefore the Root test (Theorem 8) implies the convergence of the series. The converse is not true (See Problem 5).

- (b) Follows from (a).
9. (a) Compare the Dirichlet Test with Problem 15 of the Practice Problems 11. Repeat the steps (a)-(c) given in the problem mentioned above by taking  $b_n$  in place of  $\frac{1}{n}$ .
- (b) Compare Abel's Test with Problem 15 of the Practice Problems 11. Repeat the steps (a)-(c) given in the problem mentioned above by taking  $b_n$  in place of  $\frac{1}{n}$ . In Abel's test ( $b_n$ ) could be increasing. However, the proofs of the steps (a)-(c) go through.
- (c) Apply the Dirichlet test or consider the sequence of partial sums.
- (d) Apply Abel's test.