

# FARTHEST POINTS AND THE FARTHEST DISTANCE MAP

PRADIPTA BANDYOPADHYAY AND S DUTTA

ABSTRACT. In this paper, we consider farthest points and the farthest distance map of a closed bounded set in a Banach space. We show, *inter alia*, that a strictly convex Banach space has the Mazur-like intersection property for weakly compact sets if and only if every such set is the closed convex hull of its farthest points, and recapture a classical result of Lau in a broader set-up. We obtain an expression for the subdifferential of the farthest distance map in the spirit of Preiss' Theorem which in turn extends a recent result of Westphal and Schwartz, showing that the subdifferential of the farthest distance map is the unique maximal monotone extension of a densely defined monotone operator involving the duality map and the farthest point map.

## 1. INTRODUCTION

We work with real scalars. The closed unit ball and the unit sphere of a Banach space  $X$  will be denoted by  $B(X)$  and  $S(X)$  respectively. Our notations are otherwise standard. Any unexplained terminology can be found in [3].

For a closed and bounded set  $K$  in a Banach space  $X$ , the farthest distance map  $r_K$  is defined as  $r_K(x) = \sup\{\|z - x\| : z \in K\}$ ,  $x \in X$ . For  $x \in X$ , we define the farthest point map as  $Q_K(x) = \{z \in K : \|z - x\| = r_K(x)\}$ , *i.e.*, the set of points of  $K$  farthest from  $x$ . Note that this set may be empty. Let  $D(K) = \{x \in X : Q_K(x) \neq \emptyset\}$ . The set of farthest points of  $K$  will be denoted by  $\text{far}(K)$ , *i.e.*,  $\text{far}(K) = \cup\{Q_K(x) : x \in D(K)\}$ . Call a closed and bounded set  $K$  densely remotal if  $D(K)$  is norm dense in  $X$ .

We say that a Banach space  $X$  has the Mazur Intersection Property (MIP) if every closed bounded convex set in  $X$  is the intersection of closed balls containing it. The MIP is a well studied notion in geometry of Banach space and several authors have studied Mazur-like intersection properties for different families of closed bounded convex sets. See [1, 2] and references thereof for a survey and unified treatments. However, no complete characterization is available, in particular, for every weakly compact convex set in  $X$  to be intersection of balls.

---

2000 *Mathematics Subject Classification.* Primary 46B20; Secondary 47H05.

*Key words and phrases.* Mazur Intersection Property, farthest points, densely remotal sets, farthest distance map, LUR, subdifferential.

Lau [7, Theorem 3.3] had shown that a reflexive Banach space  $X$  has the MIP if and only if every closed bounded convex set in  $X$  is the closed convex hull of its farthest points. In Section 2, we show that in a strictly convex Banach space  $X$ , every weakly compact convex set is intersection of balls if and only if every such set is the closed convex hull of its farthest points. Similar conclusions hold for compact convex sets, compact convex sets of finite affine dimension. And if  $X$  has the Radon-Nikodým Property (RNP), then similar result holds for  $w^*$ -compact convex sets in  $X^*$ .

For a closed and bounded set  $K \subseteq X$ ,  $x \in X$  and  $\alpha > 0$ , a crescent of  $K$  determined by  $x$  and  $\alpha$  is the set  $C(K, x, \alpha) = \{z \in K : \|z - x\| > r_K(x) - \alpha\}$ .

The simple but crucial observation in proving the main result in our next section is that for each of the above mentioned families of sets, every crescent of such a set contains a farthest point of the set. This also gives Lau's result in [7, Theorem 3.3] as an obvious corollary.

Recall that the subdifferential of a convex function  $\phi : X \rightarrow \mathbb{R}$  at  $x \in X$  is

$$\partial\phi(x) = \{x^* \in X^* : x^*(y - x) \leq \phi(y) - \phi(x) \text{ for all } y \in X\}.$$

The subdifferential of the function  $\phi(x) = \frac{1}{2}\|x\|^2$  is referred to as the duality map on  $X$  and is denoted by  $\mathcal{D}$ .

Since  $r_K$  is a continuous convex function,  $\partial r_K$  is a maximal monotone operator defined on  $X$ . In [9, Proposition 4.3], the authors showed that if  $X$  is a reflexive Banach space with  $X^*$  Fréchet smooth, then for a closed bounded set  $K$ ,  $\partial r_K$  is the unique maximal monotone extension of  $\mathcal{D} \frac{I - Q_K}{r_K}$  and for each  $x \in X$ ,

$$(1) \quad \partial r_K(x) = \bigcap_{\delta > 0} \overline{\text{co}}\{\mathcal{D} \frac{I - Q_K}{r_K}(y) : \|y - x\| < \delta\}.$$

Note that this is actually the Preiss' Theorem (see [8]) for  $\partial r_K$ .

For a nonreflexive space, such a statement needs qualification as  $\mathcal{D} \frac{I - Q_K}{r_K}(y)$  may be empty for some  $y$ . Nonetheless, even in nonreflexive spaces, for a densely remotal set  $K$  (e.g.,  $K$  weakly compact),  $\mathcal{D} \frac{I - Q_K}{r_K}$  is a well-defined monotone operator with dense domain. We show that if  $X$  is LUR, then  $\partial r_K$  remains the unique maximal monotone extension of  $\mathcal{D} \frac{I - Q_K}{r_K}$  and an analogue of (1) is available where we need to take the  $w^*$ -closure and choose  $y$  from the set  $D_1(K)$  defined below. We believe this provides with the only version of the Preiss' Theorem for  $\partial r_K$  is the non reflexive case.

Let  $x \in X$  and  $k \in Q_K(x)$ . We say  $x \in D_1(K)$  if  $k$  is contained in crescents of  $K$ , determined by  $x$  of arbitrarily small diameter. It is easy to note that if  $x \in D_1(K)$ , then  $Q_K(x)$  is necessarily singleton.

2. INTERSECTION OF BALLS AND FARTHEST POINTS

Here is the main theorem of this section. As mentioned in the introduction, this, in particular, gives the only known characterization of when every weakly compact convex set in  $X$  is intersection of balls.

**Theorem 2.1.** (a) *If  $X$  is a strictly convex Banach space and  $\mathcal{C}$  is one of the following families of sets,*

(i)  $\mathcal{K} = \{\text{all compact convex sets in } X\}$ .

(ii)  $\mathcal{F} = \{\text{all compact convex sets in } X \text{ with finite affine dimension}\}$ .

(iii)  $\mathcal{W} = \{\text{all weakly compact convex set in } X\}$ .

*then every  $K \in \mathcal{C}$  is intersection of balls if and only if every  $K \in \mathcal{C}$  is the closed convex hull of its farthest points.*

(b) *If  $X$  has the RNP, then  $X^*$  has the  $w^*$ -MIP if and only if every  $w^*$ -compact convex set in  $X^*$  is the  $w^*$ -closed convex hull of its farthest points.*

(c) [7] *If  $X$  is reflexive, then  $X$  has the MIP if and only if every closed bounded convex set in  $X$  is the closed convex hull of its farthest points.*

*Proof.* (a) We give the proof for the family  $\mathcal{W}$  of weakly compact sets. The same proof works in the other cases too.

*Necessity :* Let  $K \in \mathcal{W}$  and thus by [7, Theorem 2.3],  $K$  is densely remotal. We claim every crescent of  $K$  contains a farthest point of  $K$ .

To see this, let  $C(K, x, \alpha)$  be any crescent of  $K$ , Choose  $\varepsilon$  such that  $0 < \varepsilon\alpha/2$  and then take  $\beta$  such that  $0 < \beta < \alpha - 2\varepsilon$ . Since  $K$  is densely remotal, there exists  $y \in D(K)$  such that  $\|x - y\| < \varepsilon$ . Then  $C(K, y, \beta) \subseteq C(K, x, \alpha)$  and clearly  $Q_K(y) \subseteq C(K, y, \beta)$ .

Now let  $L = \overline{\text{co}}(\text{far}(K))$ . Suppose  $K \setminus L \neq \emptyset$ . Then  $L$  is the intersection of balls containing it as well and hence there exists a crescent  $C$  of  $K$  disjoint from  $L$ . By above observation  $C \cap \text{far}(K) \neq \emptyset$ . But, of course,  $\text{far}(K) \subseteq L$ . This proves the necessity.

*Sufficiency :* Suppose there exists  $K \in \mathcal{W}$  that is not intersection of balls. Let  $\tilde{K} = \bigcap \{B : B \text{ is a closed ball and } K \subseteq B\}$ . Let  $x_0 \in \tilde{K} \setminus K$ . Choose  $y_0 \in K$  and  $0 < \lambda < 1$  such that  $z_0 = \lambda x_0 + (1 - \lambda)y_0 \notin K$ .

Let  $K_1 = \text{co}(K \cup \{z_0\})$ . Then  $K_1 \in \mathcal{W}$ . We will show that  $\text{far}(K_1) \subseteq K$ , and hence,  $K_1 \neq \overline{\text{co}}(\text{far}(K_1))$ .

Let  $x \in X$ . Then  $\tilde{K} \subseteq \{u \in X : \|u - x\| \leq r_K(x)\}$ . Note that  $r_K(x) \leq r_{K_1}(x) \leq r_{\tilde{K}}(x) = r_K(x)$ . Clearly,  $z_0$  as well as any point of the form

$$(2) \quad v = \alpha z_0 + (1 - \alpha)z, \quad \alpha \in (0, 1], \quad z \in K,$$

are not extreme points of  $\tilde{K}$ , and since  $X$  is strictly convex, they are not farthest points as well. Therefore,  $\|v - x\| < r_K(x)$ . Thus,  $Q_{K_1}(x) \subseteq K$ . Since  $x \in X$  was arbitrary,  $\text{far}(K_1) \subseteq K$ .

(b) If  $X$  has the RNP, by [4, Proposition 3], each  $w^*$ -compact set  $K \subseteq X^*$  is densely remotal. Thus, necessity can be proved as in (a).

For sufficiency, note that if there exists a  $w^*$ -compact set  $K$  that is not intersection of balls, since  $K = \bigcap_{\lambda > 0} [K + \lambda B(X^*)]$ , passing to some  $K + \lambda B(X^*)$  if necessary, we may assume that  $K$  has nonempty interior. Now if we choose  $y_0 \in \text{int}(K)$ , then  $z_0$  and any point of the form (2) are interior points of  $\tilde{K}$ , and hence the result follows as before.  $\square$

The following observation is immediate from the above arguments.

**Proposition 2.2.** *Every closed bounded convex set in  $X$  is the closed convex hull of its farthest points if and only if*

- (a)  $X$  has the MIP; and
- (b) for every closed bounded convex set  $K \subseteq X$  and every closed ball  $B$  with  $K \setminus B \neq \emptyset$ ,  $K \setminus B$  contains a farthest point of  $K$ .

Note that the proof of Theorem 2.1 shows that if  $K$  is densely remotal, then  $K$  satisfies (b) above. Following example shows that we cannot dispense with (b).

**Example 2.3.** The space  $c_0$  has a strictly convex Fréchet differentiable renorming [3, Theorem 7.1 (ii)] which, thus, has the MIP. However, since the unit ball of the usual norm on  $c_0$  lacks extreme points, it must lack farthest points in the new norm.

**Remark 2.4.** This also shows that even if  $X^*$  has the RNP, there may exist a closed bounded convex set in  $X$  with  $\text{far}(K) = \emptyset$  (Compare this with [4, Proposition 3]).

Observe that since the bidual of a space with the MIP has the  $w^*$ -MIP, Theorem 2.1(b) shows that every  $w^*$ -compact convex set in  $\ell^\infty$ , with the bidual of the above norm, is the  $w^*$ -closed convex hull of its farthest points. Thus, there is a closed bounded convex set  $K \subseteq c_0$ , such that no farthest point of the  $w^*$ -closure of  $K$  in  $X^{**}$  comes from  $K$ .

### 3. THE FARTHEST DISTANCE MAP

We begin by collecting some simple properties of the set  $D_1(K)$ . Recall that a sequence  $\{z_n\} \subseteq K$  is called a maximizing sequence for  $x$  if  $\|x - z_n\| \rightarrow r_K(x)$ .

**Proposition 3.1.** *Let  $K$  be closed bounded set in a Banach space  $X$ .*

- (a)  $x \in D_1(K)$  if and only if any maximizing sequence for  $x$  converges.
- (b) If  $x \in D_1(K)$ , then  $Q_K$  is single valued and continuous at  $x$  and  $Q_K(x)$  is a strongly exposed point of  $K$ .

(c)  $D_1(K)$  is a  $G_\delta$  in  $X$ .

The following proposition shows that any discussion on  $D_1(K)$  naturally require some convexity conditions on the norm.

**Proposition 3.2.** (a) *A Banach space  $X$  is strictly convex if and only if for every compact set  $K$  and  $k \in \text{far}(K)$ , there exists  $x \in D_1(K)$ , such that  $Q_K(x) = \{k\}$ .*

(b) *A Banach space  $X$  is LUR if and only if for every closed bounded set  $K$  and  $k \in \text{far}(K)$ , there exists  $x \in D_1(K)$ , such that  $Q_K(x) = \{k\}$ .*

*Proof.* (a) Let  $K$  be a compact set in a strictly convex Banach space  $X$ . Let  $k \in \text{far}(K)$ . Then,  $k \in Q_K(x)$  for some  $x \in D(K)$ . Let  $t > 1$ . Strict convexity of the norm shows that for  $y = k + t(x - k)$ ,  $Q_K(y) = \{k\}$ . Now compactness shows that  $y \in D_1(K)$ .

Conversely, if  $X$  is not strictly convex, there exists  $x, y \in S(X)$  such that the line segment  $[x, y] \subseteq S(X)$ . Clearly,  $K = [x, y]$  is compact and  $K \subseteq Q_K(0)$ . But any point of the open segment  $(x, y)$  cannot be strongly exposed and therefore, cannot be in the set  $Q_K(D_1(K))$ .

(b) Observe that  $S(X) \subseteq Q_{B(X)}(0)$ . For any  $x \in S(X)$  and any sequence  $\{x_n\} \subseteq B(X)$  that is maximizing for  $-x$ , we have  $\|x + x_n\| \rightarrow 2$ . So if  $X$  is LUR,  $x_n \rightarrow x$ . Thus  $-x \in D_1(B(X))$  and  $x \in Q_{B(X)}(-x)$ . Now for a closed bounded set  $K \subseteq X$  and  $k \in \text{far}(K)$ , get  $x \in D(K)$  such that  $k \in Q_K(x)$ , and apply this argument with suitable translation and scaling to the ball  $B[x, r_K(x)]$ .

To prove the converse, let  $K = B(X)$ . Then,  $S(X) = \text{far}(B(X))$ . So by the hypothesis, it follows that every point in  $S(X)$  is a strongly exposed point of  $B(X)$ , and therefore,  $X$  is strictly convex.

Now let  $x_0 \in S(X)$ . By hypothesis, there exists  $x \in D_1(B(X))$  such that  $Q_K(x) = \{x_0\}$ . Then  $\|x - x_0\| = r_{B(X)}(x) = 1 + \|x\|$ . By strict convexity, it follows that  $x = \alpha x_0$  for some  $\alpha \in \mathbb{R}$  and  $|\alpha - 1| = 1 + |\alpha|$ . Therefore,  $\alpha < 0$ .

To show  $X$  is LUR, let  $\{x_n\} \subseteq B(X)$  be such that  $\|x_n + x_0\| \rightarrow 2$ . For each  $n$  consider the function on  $(0, 1)$ ,

$$f_n(\lambda) = 1 - \|\lambda x_n + (1 - \lambda)x_0\|$$

Then for all  $\lambda \in (0, 1)$ ,  $f_n(\lambda) \geq 0$ . And by triangle inequality,

$$2f_n(1/2) \geq f_n(\lambda) + f_n(1 - \lambda) \geq f_n(\lambda) \geq 0$$

By assumption,  $f_n(1/2) \rightarrow 0$ . Thus, for any  $\lambda \in (0, 1)$ ,  $f_n(\lambda) \rightarrow 0$ . In particular, putting  $\lambda = 1/(1 - \alpha)$ , we get  $\|x_n - \alpha x_0\| \rightarrow (1 - \alpha)$ , that is  $\{x_n\}$  is a maximizing sequence for  $x = \alpha x_0$ . Hence,  $x_n \rightarrow x_0$ .  $\square$

The following two lemmas are crucial in proving our main theorem of this section.

**Lemma 3.3.** *Suppose  $X$  is LUR and  $K \subseteq X$  is densely remotal. Then  $D_1(K)$  is a dense  $G_\delta$  in  $X$ .*

*Proof.* By Proposition 3.1(d), it suffices to show that  $D_1(K)$  is dense in  $X$ .

Let  $x \in D(K)$ . Get  $k \in Q_K(x)$ .  $0 < \varepsilon < 1$ . Let  $y = k + (1 + \varepsilon)(x - k)$ . Then,  $\|x - y\| = \varepsilon r_K(x)$ . It is easy to see that  $r_K(y) = (1 + \varepsilon)r_K(x)$  and by strict convexity,  $k$  is a unique farthest point from  $y$ .

We now claim  $y \in D_1(K)$ . Let  $\{z_n\} \subseteq K$  be a maximizing sequence for  $y$ . That is,  $\|z_n - y\| \rightarrow (1 + \varepsilon)r_K(x)$ . Then,

$$\left\| \frac{(z_n - x) + \varepsilon(k - x)}{(1 + \varepsilon)} \right\| \rightarrow r_K(x)$$

Then  $y_n = (z_n - x)/r_K(x) \in B(X)$ ,  $y_0 = (k - x)/r_K(x) \in S(X)$ , and for  $\lambda = 1/(1 + \varepsilon)$ , we have  $\|\lambda y_n + (1 - \lambda)y_0\| \rightarrow 1$ . Notice that since  $\varepsilon < 1$ ,  $1/2 < \lambda < 1$ . As in the proof of Proposition 3.2(b), let  $f_n(\lambda) = 1 - \|\lambda y_n + (1 - \lambda)y_0\|$ . By convexity of the norm,

$$f_n(\lambda) \geq (2 - 2\lambda)f_n(1/2) \geq 0$$

Since  $f_n(\lambda) \rightarrow 0$ , we have that  $f_n(1/2) \rightarrow 0$ , that is,  $\|y_n + y_0\| \rightarrow 2$ . Since  $X$  is LUR,  $y_n \rightarrow y_0$  and hence,  $z_n \rightarrow k$ .  $\square$

**Remark 3.4.** It follows that for any weakly compact set  $K$  in a LUR Banach space,  $D_1(K)$  is a dense  $G_\delta$  in  $X$ . So our result is more general than [5, Corollary 2.8], where it is proved that if the norm on  $X^*$  is Fréchet differentiable, then for any closed and bounded subset  $K \subseteq X$ ,  $D_1(K)$  is residual.

**Lemma 3.5.** *Let  $x \in D_1(K)$ . Then,  $\partial r_K(x) = \mathcal{D}^{\frac{I-Q_K}{r_K}}(x)$ .*

*Moreover,  $r_K$  is Gâteaux (resp. Fréchet) differentiable at  $x$  if and only if the norm is Gâteaux (resp. Fréchet) differentiable at  $x - Q_K(x)$ .*

*Proof.* Let  $Q_K(x) = \{k\}$  and  $x^* \in \mathcal{D}(\frac{x-k}{r_K(x)})$ . Then  $x^*(x - k) = r_K(x)$ . For  $z \in X$ ,  $x^*(z - x) = x^*(z) - x^*(k) - r_K(x) \leq r_K(z) - r_K(x)$ . Thus  $x^* \in \partial r_K(x)$ .

Conversely, let  $x^* \in \partial r_K(x)$ . Since  $\mathcal{D}^{\frac{I-Q_K}{r_K}}(x)$  is a  $w^*$ -closed convex subset of  $S(X^*)$ , it is enough to show that for any  $z \in S(X)$ , there is an  $x_0^* \in \mathcal{D}^{\frac{I-Q_K}{r_K}}(x)$  such that  $x^*(z) \leq x_0^*(z)$ .

Let  $\{k_n\} \subseteq K$  be such that  $\|x + z/n - k_n\| > r_K(x + z/n) - 1/n^2$ . Then  $\{k_n\}$  is a maximizing sequence for  $x$ , and hence,  $k_n \rightarrow k$ . Now

$$x^*\left(\frac{z}{n}\right) = x^*\left(x + \frac{z}{n}\right) - x^*(x) \leq r_K\left(x + \frac{z}{n}\right) - r_K(x) < \left\|x + \frac{z}{n} - k_n\right\| - r_K(x) + \frac{1}{n^2}.$$

Choose  $x_n^* \in \mathcal{D}(x + z/n - k_n)$ . Then

$$x_n^*\left(\frac{z}{n}\right) = x_n^*\left(x + \frac{z}{n} - k_n\right) - x_n^*(x - k_n) \geq \left\|x + \frac{z}{n} - k_n\right\| - r_K(x).$$

Combining the two, we have  $x^*(z) \leq x_n^*(z) + 1/n$ . Let  $x_0^*$  be a  $w^*$ -cluster point of  $\{x_n^*\}$ . Since  $x + z/n - k_n$  converges to  $x - k$  in norm, we have  $x_0^* \in \mathcal{D}_{\frac{I-Q_K}{r_K}}(x)$  and  $x^*(z) \leq x_0^*(z)$ , as desired.

Thus, the norm is Gâteaux differentiable at  $x - k \Leftrightarrow \mathcal{D}(\frac{x-k}{r_K(x)})$  is singleton  $\Leftrightarrow$  so is  $\partial r_K(x) \Leftrightarrow r_K$  is Gâteaux differentiable at  $x$ .

Now, let  $\{x^*\} = \partial r_K(x) = \mathcal{D}(\frac{x-k}{r_K(x)})$ . For any  $\lambda \in \mathbb{R}$  and  $z \in B(X)$ ,  $x^*(\lambda z) \leq \|x + \lambda z - k\| - \|x - k\| \leq r_K(x + \lambda z) - r_K(x)$ . Therefore,

$$\left| \frac{\|x + \lambda z - k\| - \|x - k\|}{\lambda} - x^*(z) \right| \leq \left| \frac{r_K(x + \lambda z) - r_K(x)}{\lambda} - x^*(z) \right|.$$

Thus, Fréchet differentiability of  $r_K$  at  $x$  implies that of the norm at  $x - k$ .

Conversely, let the norm be Fréchet differentiable at  $x - k$ . Let  $x_n \rightarrow x$ ,  $x_n^* \in \partial r_K(x_n)$  and  $x^* \in \partial r_K(x)$ , then  $\{x_n^*\} \subseteq B(X^*)$  and since  $r_K$  is Gâteaux differentiable at  $x$ ,  $x_n^* \rightarrow x^*$  in the  $w^*$ -topology. Since  $x^* \in \mathcal{D}_{\frac{I-Q_K}{r_K}}(x)$ ,  $x^*$  is a  $w^*$ -norm point of continuity of  $B(X^*)$ , and therefore,  $x_n^* \rightarrow x^*$  in norm. It follows that  $r_K$  is Fréchet differentiable at  $x$ .  $\square$

**Remark 3.6.** [5, Theorem 3.2(a)] proves only the “necessity” part of this result. Our proof is also simpler.

Combining Lemma 3.5 with Lemma 3.3, it follows that in a Banach space with smooth LUR norm, the farthest distance map  $r_K$  of a densely remotal set  $K$  is Gâteaux differentiable on a dense  $G_\delta$ .

We now state the main theorem of this section. This gives an expression for  $\partial r_K$  in the spirit of Preiss’ Theorem [8]. Note that our result does not need smoothness of the norm and with (Fréchet) smoothness, by Theorem 3.5, we get back Preiss’ Theorem for  $\partial r_K$ .

**Theorem 3.7.** *Let  $K$  be such that  $D_1(K)$  is dense in  $X$  and  $x \in X$ . Then*

$$\partial r_K(x) = \bigcap_{\delta > 0} \overline{\text{co}}^* \{ \partial r_K(y) : y \in D_1(K) \text{ and } \|y - x\| < \delta \}.$$

*Proof.* Let  $x^* \in RHS$  and  $\varepsilon > 0$ . Choose  $\delta < \varepsilon/3$ . For  $z \in X$ , choose  $y \in D_1(K)$  and  $y^* \in \partial r_K(y)$  such that  $\|y - x\| < \delta$  and  $x^*(z - x) < y^*(z - x) + \delta$ . Thus,

$$\begin{aligned} x^*(z - x) &< y^*(z - x) + \delta = y^*(z - y) + y^*(y - x) + \delta \leq r_K(z) - r_K(y) + 2\delta \\ &\leq r_K(z) - r_K(x) + 3\delta \leq r_K(z) - r_K(x) + \varepsilon. \end{aligned}$$

Since  $\varepsilon$  is arbitrary, we have  $x^* \in \partial r_K(x)$ .

Conversely, let  $x^* \in \partial r_K(x)$ . As in Lemma 3.5, we will show given any  $z \in S(X)$  there is an  $x_0^* \in RHS$  such that  $x^*(z) \leq x_0^*(z)$ .

For each  $n$ , get  $y_n \in D_1(K)$  such that  $\|x + z/n - y_n\| < 1/n^2$ . Then

$$x^*\left(\frac{z}{n}\right) \leq r_K\left(x + \frac{z}{n}\right) - r_K(x) \leq r_K(y_n) - r_K\left(y_n - \frac{z}{n}\right) + \frac{2}{n^2}.$$

Let  $x_n^* \in \partial r_K(y_n)$ . Let  $k_n \in Q_K(y_n)$ . Then

$$x_n^*\left(\frac{z}{n}\right) = x_n^*(y_n - k_n) - x_n^*\left(y_n - \frac{z}{n} - k_n\right) \geq r_K(y_n) - r_K\left(y_n - \frac{z}{n}\right).$$

Thus  $x^*(z) \leq x_n^*(z) + 2/n$ . Let  $x_0^*$  be a  $w^*$ -cluster point of  $x_n^*$ . Then  $x_0^* \in RHS$  and  $x^*(z) \leq x_0^*(z)$ .  $\square$

Combining the Lemma 3.3, Lemma 3.5 and Theorem 3.7 we obtain the following:

**Corollary 3.8.** *Suppose  $K$  is a densely remotal set in a LUR Banach space  $X$ . Then  $\partial r_K$  is the unique maximal monotone extension of the densely defined monotone operator  $\mathcal{D}\frac{I-Q_K}{r_K}$  and for each  $x \in X$ , we have,*

$$\partial r_K(x) = \bigcap_{\delta > 0} \overline{co}^* \left\{ \mathcal{D}\frac{I-Q_K}{r_K}(y) : y \in D_1(K) \text{ and } \|y - x\| < \delta \right\}.$$

**Remark 3.9.** In [9, Proposition 4.3] obtained the similar conclusion for reflexive Banach spaces with  $X^*$  Fréchet smooth.

We end this section with a result on range of  $\partial r_K$ . Compare this with [9, Theorem 4.2].

**Theorem 3.10.** *Let  $X$  be a smooth (resp. Fréchet smooth) Banach space. Let  $K \subseteq X$  be a closed and bounded set such that  $D_1(K)$  is dense in  $X$ , then the image of  $D_1(K)$  under  $\partial r_K$  is  $w^*$ -dense (resp. norm dense) in  $S(X^*)$ .*

*Proof.* Let  $NA(X)$  denote the set of norm attaining functionals in  $S(X^*)$ . By Bishop-Phelps Theorem,  $NA(X)$  is norm dense in  $S(X^*)$ . Let  $x_0^* \in NA(X)$  and  $x_0 \in S(X)$  such that  $x_0^*(x_0) = 1$ . By density of  $D_1(K)$ , choose  $x_n \in D_1(K)$  such that  $\|x_n - nx_0\| < 1/n$  and let  $x_n^* \in \partial r_K(x_n)$ . Then  $\|x_n\| \rightarrow \infty$ . Therefore, by [9, Lemma 4.1],  $\lim x_n^*(x_n/\|x_n\|) = \lim \|x_n^*\| = 1$ . But since  $x_n/\|x_n\| \rightarrow x_0$  in norm,  $x_n^*(x_0) \rightarrow 1$  as well. Thus, any  $w^*$ -cluster point of  $\{x_n^*\}$  is in  $\mathcal{D}(x_0)$ . Since the norm is smooth, this set is singleton. Hence,  $x_n^* \rightarrow x_0^*$  in  $w^*$ -topology.

Now, if the norm on  $X$  is Fréchet smooth, then  $x_0^*$  chosen above is a  $w^*$ -norm point of continuity of  $B(X^*)$ . Thus  $\partial r_K(D_1(K))$  is norm dense in  $S(X^*)$ .  $\square$

## REFERENCES

- [1] P. Bandyopadhyay, *The Mazur Intersection Property for Families of Closed Bounded Convex Sets in Banach Spaces*, Colloq. Math. **LXIII** (1992), 45–56.
- [2] D. Chen and B. L. Lin, *Ball separation properties in Banach spaces*, Rocky Mountain J. Math., **28** (1998), 835–873.
- [3] R. Deville, G. Godefroy, and V. Zizler, *Smoothness and renormings in Banach spaces*, Pitman Monographs and Surveys in Pure and Applied Mathematics, 64. Longman Scientific & Technical, Harlow; 1993.
- [4] R. Deville and V. Zizler, *Farthest Points in  $W^*$ -Compact Sets*, Bull. Austral. Math. Soc. **38** (1988), 433–439.



- [5] S. Fitzpatrick, *Metric Projections and the Differentiability of Distance Functions*, Bull. Austral. Math. Soc. **22** (1980), 291–312.
- [6] J. R. Giles, D. A. Gregory and B. Sims, *Characterisation of Normed Linear Spaces with the Mazur's Intersection Property*, Bull. Austral. Math. Soc. **18** (1978), 105–123.
- [7] K.-S. Lau, *Farthest Points in Weakly Compact Sets*, Israel J. Math. **22** (1975), 168–174.
- [8] D. Preiss, *Fréchet Derivatives of Lipschitz functions*, J. Funct. Anal. **91** (1990), 312–345.
- [9] U. Westphal and T. Schwartz, *Farthest Points and Monotone Operators*, Bull. Austral. Math. Soc. **58** (1998), 75–92.

(Pradipta Bandyopadhyay) STAT–MATH DIVISION, INDIAN STATISTICAL INSTITUTE, 203, B. T. ROAD, KOLKATA 700 108, INDIA, *E-mail* : [pradipta@isical.ac.in](mailto:pradipta@isical.ac.in)

(S Dutta) STAT–MATH DIVISION, INDIAN STATISTICAL INSTITUTE, 203, B. T. ROAD, KOLKATA 700 108, INDIA, *E-mail* : [sudipta\\_r@isical.ac.in](mailto:sudipta_r@isical.ac.in)