

## Lecture 7

### **Cauchy Theorem:**

Let  $f$  be analytic inside and on a simple, closed, piecewise smooth curve  $C$ . Then,

$$\int_C f(z) dz = 0.$$

**Definitions:** Let  $z(t)$ ,  $a \leq t \leq b$ , be parametric representation of the curve  $C$ .

**Simple Curve:** The curve  $C$  is said to be **simple**, if it does not have any self-intersections

(i.e.  $z(t_1) \neq z(t_2)$  whenever  $t_1 \neq t_2$  ( $a < t_1, t_2 < b$ )).

**Closed Curve:** The curve  $C$  is said to be **Closed**, if end point of the curve is the same as its initial point

(i.e.  $z(a) = z(b)$ ).

**Piece-wise smooth Curve:** The curve  $C$  is said to be **Piece-wise smooth**, if  $z(t)$  is piece-wise differentiable (i.e. differentiable for

all except finitely many  $t$ ) and  $\frac{d}{dt} z(t)$  (denoted as  $\dot{z}(t)$ ) is piece-

wise continuous in the interval  $[a, b]$

**Proof (Under the assumption that  $f'(z)$  is continuous on  $C$ )**

By Green's Theorem,

$$\int_C Pdx + Qdy = \iint_R (Q_x - P_y) dx dy ,$$

where, curve  $C$  is boundary of the region  $R$  and the first partial derivatives  $P, Q, Q_x, P_y$  exist and are continuous in  $C \cup R$ .

The hypothesis of Cauchy Theorem implies that the conditions of Green's Theorem are satisfied.


$$\text{Now, } \int_C f(z) dz = \int_a^b f(z(t)) \dot{z}(t) dt$$

$$= \int_a^b (u + iv)(\dot{x}(t) + i\dot{y}(t)) dt$$

$$= \int_a^b (u\dot{x} - v\dot{y}) dt + i \int_a^b (u\dot{y} + v\dot{x}) dt$$

$$= \int_C u dx - v dy + i \int_C u dy + v dx$$

$$= -\iint_R (u_y + v_x) dx dy + i \iint_R (u_x - v_y) dx dy$$




The proof of Cauchy Theorem in the general case, where the continuity of  $f'(z)$  is not assumed, is beyond the scope of this course.


**Cauchy Theorem for Multiply Connected Domains** (Domain with holes).

Let simple closed piece-wise smooth curves  $C_1, \dots, C_n$  be enclosed by a simple, closed piece-wise smooth curve  $C$ , all the curves being oriented anticlockwise. Let  $D$  be domain with boundary curves  $C, C_1, \dots, C_n$  (**Such a domain is called a multiply connected domain**). If a function  $f(z)$  is analytic on  $D \cup C \cup C_1 \cup \dots \cup C_n$ , then

$$\oint_C f(z) dz = \oint_{C_1} f(z) dz + \dots + \oint_{C_n} f(z) dz.$$



**Proof:** Join  $C$  (oriented anticlockwise) and  $C_1^*, \dots, C_n^*$  (the curves  $C_1, \dots, C_n$  oriented clockwise) by straight line segment as shown in the figure for  $n = 3$ . Observe that with these orientations,  $D$  lies to left if one traverses along any of these curves.



Applying Cauchy Theorem to the simply connected domain bounded by the curve

$$\Gamma = L_1 \cup C_1^{*u} \cup L_2 \cup C_2^{*u} \cup L_3 \cup C_3^{*u} \cup \dots \cup L_n \cup C_n^* \cup \\ -L_n \cup C_{n-1}^{*l} \cup \dots -L_3 \cup C_2^{*l} \cup -L_2 \cup C_1^{*l} -L_1 \cup C$$

where,  $C_i^{*u}$  denotes the upper part of the curve  $C_i^*$  and  $C_i^{*l}$  denotes the lower part of the curve  $C_i^*$  (observe that  $\Gamma$  has positive orientation, since the domain bounded by it lies to its left when one traverses on  $\Gamma$ ), it follows that

$$\oint_C f(z) dz + \oint_{C_1^*} f(z) dz + \dots + \oint_{C_n^*} f(z) dz = 0$$

(since the integrals along  $L_i$ 's are equal and opposite to each other)

$$\begin{aligned} \Rightarrow \oint_C f(z) dz &= \oint_{-C_1^*} f(z) dz + \dots + \oint_{-C_n^*} f(z) dz \\ &= \oint_{C_1} f(z) dz + \dots + \oint_{C_n} f(z) dz \end{aligned}$$

**Corollary.** If  $f$  is analytic (i) on two simple, closed, piece-wise smooth curves  $C_1$  and  $C_2$  and (ii) inside the domain bounded by  $C_1$  and  $C_2$ , then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz.$$

The above corollary helps in evaluation of integrals over curve  $C_1$ , the determination of whose parametric representation may be complicated. In such a case, the possibility of obtaining a curve  $C_2$  satisfying the conditions of the corollary and whose parametric representation is simple to obtain, is explored and the integral is evaluated with the help of above corollary.

**Example:** Evaluate  $\oint_{\Gamma} \frac{1}{w - z_0} dw$ , where  $\Gamma$  is any anticlockwise oriented simple closed piecewise smooth curve and  $z_0$  is a point lying in the bounded domain  $D$  with boundary  $\Gamma$ .

Note that direct evaluation of the above integral is not possible, since any explicit equation of  $\Gamma$  is not known. However, this integral could be simply evaluated by using the above theorem.

Consider any anticlockwise oriented circle  $C_r : |w - z_0| = r$ , with  $r$  small enough so that  $C_r$  lies in  $D$ . The function  $\frac{1}{w - z_0}$  is analytic on the curves  $\Gamma$  and  $C_r$  and in the domain bounded by these curves. Therefore, by Cauchy Theorem for Multiply connected domains,

$$\oint_{\Gamma} \frac{1}{w - z_0} dw = \oint_{C_r} \frac{1}{w - z_0} dw = \int_0^{2\pi} \frac{1}{re^{it}} ire^{it} dt = 2\pi i$$

since,  $w(t) = z_0 + re^{it}$ ,  $0 \leq t \leq 2\pi$ , is a parametric representation of the circle  $C_r$ .

**Cauchy Integral Formula:** If  $f$  is analytic in a domain  $G$  and  $\overline{B(a,r)} \subseteq G$ , where  $\overline{B(a,r)} = \{w : |w-a| \leq r\}$ . Then, for any  $z \in \{|w-a| < r\}$

$$f(z) = \frac{1}{2\pi i} \int_{C_r} \frac{f(w)}{w-z} dw \quad (1)$$

where,  $C_r : w(t) = z + re^{it}, 0 \leq t \leq 2\pi$ .

**Proof:** Consider a circle  $|w-z| = \delta^*$  centered at  $z$  and having radius  $\delta^*$  sufficiently small such  $\{|w-z| = \delta^*\} \subset \{|w-a| < r\}$ . Then, by Cauchy Theorem of Multiply Connected Domains,

$$\int_{C_r} \frac{f(w)}{w-z} dw = \int_{|w-z|=\delta^*} \frac{f(w)}{w-z} dw$$

since the integrand is an analytic function in the domain lying between  $C_r$  and  $|w-z| = \delta^*$ . Now, note that

$$\int_{|w-z|=\delta^*} \frac{f(w)}{w-z} dw = \int_{|w-z|=\delta^*} \frac{f(w) - f(z)}{w-z} dw + f(z) \int_{|w-z|=\delta^*} \frac{1}{w-z} dw \quad (*)$$

The second term of (\*) =  $2\pi i f(z)$ . Therefore, Cauchy Integral Formula follows if we prove that the first term of (\*) is zero.

For this use continuity of  $f(w)$  at ' $z$ ', which gives that for every  $\varepsilon > 0$ , there exists  $\delta > 0$  such that

$|f(w) - f(z)| < \varepsilon$  whenever  $|w-z| < \delta$ . Choose  $\delta^* < \delta$ .



Then,

$$\left| \int_{|w-z|=\delta^*} \frac{f(w) - f(z)}{(w-z)} dw \right| < \frac{\varepsilon}{\delta^*} \times 2\pi\delta^* = 2\pi\varepsilon \text{ (by ML-Estimate)}$$

$$\Rightarrow \int_{|w-z|=\delta^*} \frac{f(w) - f(z)}{(w-z)} dw = 0 \text{ since } \varepsilon \text{ is arbitrary.}$$

**Note:** In view of Cauchy Theorem for multiply connected domains, Cauchy Integral Formula (1) remains valid with  $C_r$  replaced by any simple closed piece-wise smooth curve  $\Gamma$  so that (i) every point enclosed by  $\Gamma$  is in  $D$  (ii)  $\Gamma$  encloses the point  $z$ . This is because the function  $f(w)/(w-z)$  is analytic in the domain lying between  $C_r$  and  $\Gamma$ .