


**Note:** The problems assigned are from the text book *Introduction to Electrodynamics* (4<sup>th</sup> edition) by David J. Griffiths

1. Problem 3.32
2. Problem 3.35
3. Problem 3.43
4. Problem 3.58
5. Problem 4.6
6. Problem 4.9
7. Problem 4.11
8. Problem 4.13
9. Problem 4.15
10. Problem 4.16
11. Problem 4.18
12. Problem 4.30
13. Problem 4.31
14. Problem 4.33
15. Problem 4.35


**Hints:**

- 3.35: Net charge is zero so the first non-zero contribution to the field comes from the dipole. Dipole moment will be in the z-direction.
- 3.43: The net field inside the conductor is zero. Surface charge density  $\sigma(\theta) = k\cos\theta$  over a spherical shell gives constant field inside the sphere and a dipole field outside.
- 3.58: Use image problem for a grounded sphere and  $\sigma = -\hat{n} \cdot \vec{\nabla}V$  on a conducting surface.
- 4.6: A perfect dipole  $\vec{p} = q\vec{a}$  with  $q \rightarrow \infty$  and  $a \rightarrow 0$  or also a spherical shell with surface charge density  $\sigma(\theta) = k\cos\theta$ . Method of images can be used here to find the net field.
- 4.11: Find bound charges in each case and calculate the field given by them.
- 4.13: Two-dimensional analogue of uniformly polarized sphere.
- 4.15 and 4.16: Look at the bound charges and also at the boundary conditions.
- 4.18: Displacement remains the same throughout the dielectric.
- 4.30: Draw the field lines and look at the formula  $\vec{F} = (\vec{p} \cdot \vec{\nabla})\vec{E}$
- 4.31: It will be helpful to use planar polar coordinates here. The dipole experiences both a force and a torque.
- 4.35: Use Poisson's equation and boundary conditions at the spherical surface.


3.32



(a)



(b)



(c)

(i) Monopole is  $3q - q = 2q$  in all the cases

(ii) Dipole in

(a)	$3qa \hat{z}$	} $\vec{p} = \sum q_i \vec{a}_i$
(b)	$3qa \hat{z}$	
(c)	$3qa \hat{y}$	

(iii) Approximate potential in

$$(a) \quad \frac{1}{4\pi\epsilon_0} \left[ \frac{2q}{r} + \frac{3qa \cos\theta}{r^2} \right]$$



$$(b) \quad \frac{1}{4\pi\epsilon_0} \left[ \frac{2q}{r} + \frac{3qa \cos\theta}{r^2} \right]$$

$$(c) \quad \frac{1}{4\pi\epsilon_0} \left[ \frac{2q}{r} + \frac{3qa \sin\theta \sin\phi}{r^2} \right]$$

$$V(\vec{r}) = \frac{1}{4\pi\epsilon_0} \left[ \frac{Q}{r} + \frac{\vec{p} \cdot \hat{r}}{r^2} \right] \text{ up to dipole}$$

3.35

(2)



The red and green pair in the right sphere

$$q_{rs} \quad d\vec{p} = (\rho \cdot r^2 dr \cdot d(\cos\theta) d\phi) 2r \cos\theta \hat{z}$$

(This has taken into account the charge in the lower hemisphere too)

$$\text{So } \vec{p} = 2\rho \int_0^{2\pi} d\phi \int_0^1 \cos\theta d(\cos\theta) \int_0^R r^3 dr$$

This integral is only over the upper hemisphere because of the comment above

$$= 4\pi\rho \times \frac{R^4}{4} \times \frac{1}{2} = \left( \frac{\pi\rho R^4}{2} \right)$$


NOTE: The same answer can also be obtained by (distance of the two Centre of charges)  $\times$  (charge in each hemisphere) because this calculation is like the Centre of mass calculation.

CM for hemisphere =  $\left( \frac{3R}{8} \right)$  above the centre of sphere

$$\Rightarrow \vec{p} = \frac{2\pi R^3}{3} \rho \times 2 \times \frac{3R}{8} = \frac{\pi R^4 \rho}{2}$$

3.43

$$\sigma(\theta) = k \cos \theta$$



Potential everywhere will have three components:

(i) Potential due to the boundary condition that

$$V(a) = V_0$$

(ii) Potential due to the surface charge density on the outer shell

(iii) Potential due to the induced charge

The problem can be solved in two ways :

(a) directly by realizing that field given by

$$\sigma(\theta) = k \cos \theta \text{ inside the shell is } -\frac{k}{3\epsilon_0} \hat{z}$$

And field inside the ~~shell~~ conducting sphere is zero, and


(b) by the image method.

### DIRECT METHOD :

To make field vanish inside the conducting sphere, charge  $-k \cos \theta$  will get induced on the conducting sphere. Thus ~~after~~ after the charges are induced, the picture is as follows



(4)



Also the dipole moment

$\int \sigma(b) = k\epsilon_0 0$  is

$$\vec{p} = \frac{4\pi}{3} k R^3 \hat{z}$$

if the surface charge density

is spread over a sphere of radius  $R$

Then for

$r > b$  : the potential is the sum of  
of two potentials - that due to  $\vec{p}$  of  $\sigma = k\epsilon_0 0$   
over sphere of radius  $b$  and the other due to  
 $\vec{p}$  of  $\sigma = -k\epsilon_0 0$  over sphere of radius  $a$ .

In addition there is background potential

of  $\left(\frac{V_0 a}{r}\right)$  because  $V = V_0$  at  $r = a$


$$\text{Then } V(r > b) = \frac{1}{4\pi\epsilon_0} \left[ \frac{\frac{4\pi}{3} k b^3 \hat{z} \cdot \hat{r}}{r^2} - \frac{\frac{4\pi}{3} k a^3 \hat{z} \cdot \hat{r}}{r^2} \right] + \frac{V_0 a}{r}$$

$$= V_0 \frac{a}{r} + \frac{(b^3 - a^3)}{r^2} \frac{k\epsilon_0 0}{3\epsilon_0}$$

For  $a \leq r \leq b$ , the potential is (5)

$$\begin{aligned}
 & \frac{V_0 a}{r} + \text{potential for electric field } -\frac{k}{3\epsilon_0} \hat{z} \\
 & + \text{potential of dipole } \vec{p} = -\frac{4\pi}{3} k a^3 \hat{z} \\
 \\ 
 & = \frac{V_0 a}{r} + \frac{k z}{3\epsilon_0} - \frac{1}{4\pi\epsilon_0} \cdot \frac{4\pi}{3} \frac{k a^3 \hat{z} \cdot \hat{r}}{r^2} \\
 \\ 
 & = \frac{V_0 a}{r} + \frac{k r \cos\theta}{3\epsilon_0} - \frac{k a^3 \cos\theta}{3\epsilon_0 r^2}
 \end{aligned}$$

### IMAGE METHOD:



For the charge contained in solid angle  $d\Omega$  (see figure) there is an image inside the conductor

$$\text{Charge } dq = (d\Omega b^2) \sigma(\theta)$$

$$\text{Image } dq' = -(d\Omega b^2) \sigma(\theta) \frac{a}{b}$$

$$\sigma'(\theta) = \frac{dq'}{d\Omega \left(\frac{a^2}{b}\right)^2} = -\sigma(\theta) \frac{b^3}{a^3}$$

Radius of the image sphere

$$= -\frac{k b^3}{a^3} \cos\theta$$

(6)

Field for  $r \geq a$  by the induced charge will include field due to this image


But this image charge also is equivalent

$$\text{to a dipole} = \frac{4\pi}{3} \left(\frac{a^2}{b}\right)^3 \times \left(-k \frac{b^3}{a^3}\right) \hat{z}$$

$$= -\frac{4\pi}{3} a^3 k \hat{z}$$

This is the same dipole moment as obtained earlier and leads to the same answers.


3.58



We wish to find a  $\sigma(\theta)$  that gives the same field outside that is given by 'q' shown above.


For this consider the following situation.

(7)



Field outside  
is that given by  
charge  $q$

$\sigma(0) = 0$  by Gauss' law



Field outside is that given by the sum of  
charge  $q$  and its image  $-\frac{qR}{a}$  at  $\frac{R^2}{a}$  distance  
OR

Field outside is that given by the sum of field  
given by  $\sigma(0)$  and  $-\frac{qR}{a}$  at  $R^2/a$

$\Rightarrow$  Field given by  $q$  and  $\sigma(0)$  are the same


$$\Rightarrow \sigma(0) = -\epsilon_0 \hat{n} \cdot \nabla V = -\epsilon_0 \frac{\partial}{\partial r} \left[ \frac{q}{|\vec{r} - a\hat{z}|} - \frac{qR/a}{|\vec{r} - \frac{R^2}{a}\hat{z}|} \right]_{r=R}$$

= Answer



4.9

(8)



Let  $\vec{p}$  be in the  $xy$  plane

$$\vec{p} = p \cos \theta \hat{r} + p \sin \theta \hat{\theta}$$

$\vec{E}$  at 'q' due to the dipole

$$\vec{E} = \frac{1}{4\pi\epsilon_0} \left[ \frac{3(\vec{p} \cdot \hat{r})\hat{r} - \vec{p}}{r^3} \right]$$

Although direction of  $\vec{r}$  from  $\vec{p}$  to  $q$  is opposite to  $\hat{r}$ ,  $3(\vec{p} \cdot \hat{r})\hat{r}$  term remains unchanged because of two '-' signs occurring there

$$\begin{aligned} \text{Force on } q &= q \vec{E} = \frac{q}{4\pi\epsilon_0} \left[ \frac{3p \cos \theta \hat{r} - p \cos \theta \hat{r} - p \sin \theta \hat{\theta}}{r^3} \right] \\ &= \frac{q}{4\pi\epsilon_0} \left[ \frac{2p \cos \theta \hat{r} - p \sin \theta \hat{\theta}}{r^3} \right] \end{aligned}$$

$$\text{Force on dipole} = (\vec{p} \cdot \vec{\nabla}) \vec{E} \text{ due to } q$$


$$\begin{aligned} \text{Use } \frac{\partial}{\partial r} \hat{r} &= 0 \\ \frac{\partial}{\partial \theta} \hat{r} &= \hat{\phi} = \hat{\theta} \\ &= \left( p r \frac{\partial}{\partial r} + p \frac{1}{r} \frac{\partial}{\partial \theta} \right) \frac{kq}{r^2} \hat{r} \\ &= \left( p \cos \theta \frac{\partial}{\partial r} + \frac{p \sin \theta}{r} \frac{\partial}{\partial \theta} \right) \frac{kq}{r^2} \hat{r} \\ &= - \text{the force above} \end{aligned}$$




4.11

Done in Lecture

4.13



The charge distribution above can be generated by displacing two long cylinders carrying charge densities  $\rho$  and  $-\rho$  with respect to each other



Charge in the shaded area (blue shaded area)

$$= l \times R d\theta \cdot \cos \theta \rho$$

for length \$l\$ of the cylinder

$$dq = (R l d\theta) \rho \cos \theta$$

$$\sigma(\theta) = \frac{dq}{(R l d\theta)} = \rho \cos \theta$$

(10)

$Rl d\theta$  is the surface area along the length of the cylinder

Let  $f \rightarrow \infty$  and  $a \rightarrow 0$  such that  
 $fa \rightarrow \sigma_0$  (or  $\rho$ )


$$\sigma(\omega) = \sigma_0 \cos\theta = \rho \cos\theta$$

So field inside will be (by Gauss' law)

$$= -\frac{\rho}{2\epsilon_0} \hat{z}$$

Field outside will be superposition of fields due to two long wires carrying charge per unit length  $\lambda = \pm \pi R^2 \rho$  and displaced with respect to each other by 'a'

Field outside the cylinders



$$\vec{E}(s) = -\frac{\lambda \vec{s}}{2\pi\epsilon_0 s^2} + \frac{\lambda (\vec{s} - a\hat{x})}{2\pi\epsilon_0 |\vec{s} - a\hat{x}|^2}$$

$$\lambda = \pi R^2 \rho$$

$$\vec{E}(\vec{r}) = -\frac{\lambda \vec{r}}{2\pi\epsilon_0 s^2} + \frac{\lambda}{2\pi\epsilon_0} \frac{(\vec{r} - a\hat{x})}{(s^2 + a^2 - 2sa\cos\theta)}$$

$$= -\frac{\lambda \vec{r}}{2\pi\epsilon_0 s^2} + \frac{\lambda}{2\pi\epsilon_0} \frac{(\vec{r} - a\hat{x})}{s^2} \left[ 1 + \frac{a^2}{s^2} - \frac{2a}{s} \cos\theta \right]^{-1}$$

$$= \cancel{-\frac{\lambda \vec{r}}{2\pi\epsilon_0 s^2}} + \cancel{\frac{\lambda \vec{r}}{2\pi\epsilon_0 s^2}} + \frac{\lambda}{2\pi\epsilon_0} \frac{2a\cos\theta}{s^3} \vec{r}$$

$$- \frac{\lambda}{2\pi\epsilon_0} \frac{a\hat{x}}{s^2}$$

(keeping terms up to order 'a' in the expansion of  $\left[ 1 + \frac{a^2}{s^2} - \frac{2a}{s} \cos\theta \right]^{-1}$ )

$$= \frac{1}{2\pi\epsilon_0} \pi R^2 \left[ 2 \rho a \cos\theta \frac{\vec{r}}{s^3} - \frac{\rho a \hat{x}}{s^2} \right]$$

using  $\lambda = \pi R^2 \rho$

Now  $\rho a \rightarrow P$        $\rho \hat{x} = \vec{P}$

$$P \cos\theta = \vec{P} \cdot \hat{s} \quad \frac{\vec{r}}{s^3} = \frac{\hat{s}}{s^2}$$

$$\Rightarrow \vec{E}(\vec{r}) = \frac{R^2}{2\epsilon_0} \left[ \frac{2(\vec{P} \cdot \hat{s}) \hat{s}}{s^2} - \vec{P} \right]$$




4.15 — Done in Lecture

(12)

4.16 Cavity can be thought of as the original bulk material + material of cavity's shape with  $-\vec{P}$   
Now use principle of superposition.

4.18



$$\vec{\nabla} \cdot \vec{D}' = \rho_{\text{free}} ; \text{ By symmetry } \vec{\nabla} \times \vec{D}' = 0$$

because  $\vec{D}' \perp \text{ plates}$  (like  $\vec{E}$  without dielectrics)

$$(a) \quad \Rightarrow \quad \vec{D}' = -\sigma \hat{z}$$

$D_{\perp}$  is continuous so  $D$  is going to be the same in both slabs. ( $E = D/\epsilon_0 K$ )

$$(b) \quad E_{\text{upper slab}} = -\frac{\sigma}{2\epsilon_0} \hat{z}, \quad E_{\text{lower slab}} = -\frac{2\sigma}{3\epsilon_0} \hat{z}$$

$$(c) \quad \vec{P} = \vec{D}' - \epsilon_0 \vec{E}'$$

$$\vec{P}_{\text{upper slab}} = \left(-\sigma + \frac{\sigma}{2}\right) \hat{z} = -\frac{\sigma}{2} \hat{z}$$

$$\vec{P}_{\text{lower slab}} = \left(-\sigma + \frac{2\sigma}{3}\right) \hat{z} = -\frac{\sigma}{3} \hat{z}$$

$$(d) \quad \Delta V = \mathcal{E}_{\text{upper}} \times a + \mathcal{E}_{\text{lower}} \times a$$

$$= \frac{2\sigma}{\epsilon_0} \left( \frac{1}{2} + \frac{2}{3} \right)$$

$$= \frac{7\sigma}{6\epsilon_0}$$

$$(e) \quad \sigma_b = \vec{P} \cdot \hat{n} \quad (\oint_b = \vec{\nabla} \cdot \vec{P} = 0)$$

Diagram for upper slab:

Top surface:  $\sigma_{\text{up}} = -\sigma/2$ ,  $\uparrow \hat{n}$

Bottom surface:  $\downarrow \hat{n}$

Electric field  $\vec{P}$  is shown as downward arrows.

Calculation:  $\sigma_{\text{up}} = \vec{P} \cdot \hat{n} = \frac{\sigma}{2}$

Diagram for lower slab:

Top surface:  $\uparrow \hat{n}$ ,  $\sigma_{\text{low}} = -\sigma/3$

Bottom surface:  $\downarrow \hat{n}$


Electric field  $\vec{P}$  is shown as downward arrows.

Calculation:  $\sigma_{\text{low}} = \vec{P} \cdot \hat{n} = \frac{\sigma}{3}$

(f) Use standard sheet of charge formula

4.30

(14)




Field lines are going to be as shown  
( $\perp$  to the equipotential plates)


$$\vec{F}_{\text{on dipole}} = (\vec{p} \cdot \vec{\nabla}) \vec{E}$$

$$= p \frac{\partial}{\partial y} \vec{E} \quad \vec{p} = p \hat{y}$$

As  $y$  changes through the  $\vec{p}$ 's position  
( $x$  axis)  $\vec{E}$  changes direction from



$$\vec{E}_2 - \vec{E}_1 =$$




So  $\vec{p}$  will experience a force pointing in  
the  $+\hat{x}$  direction

4.31

Done in Lecture

(15)

4.33



$$\vec{P} = k\vec{r}$$

$$\rho_b = -\vec{\nabla} \cdot \vec{P} = -3k$$

$$\sigma_b = \vec{P} \cdot \hat{n} = + k\vec{r} \cdot \hat{x} \Big|_{x=a/2}$$

$$= + kx \Big|_{a/2}$$

$$= + \frac{ka}{2} \text{ on the surfaces}$$

$\perp$  to the  $x$ -axis

The same answer for faces  $\perp$  to  $y$ - and  $z$ -axis

$$\text{Net charge} = 6 \cdot \frac{ka}{2} \cdot a^2 - 3ka^3 = 0$$

4.35 : Done in Lecture