

Practice Problems 8: Mean Value Theorem, Cauchy Mean Value Theorem, L'Hospital's Rule

---

1. Establish the following inequalities using the MVT.

(a)  $\frac{1}{2\sqrt{n+1}} < \sqrt{n+1} - \sqrt{n} < \frac{1}{2\sqrt{n}}$  for all  $n \in \mathbb{N}$ .

(b)  $\frac{x-1}{x} < \log x < x-1$  for  $x > 1$ .

(c)  $e^x \geq ex$  for all  $x \in \mathbb{R}$ .

2. Does there exist a differentiable function  $f : [0, 2] \rightarrow \mathbb{R}$  satisfying  $f(0) = -1, f(2) = 4$  and  $f'(x) \leq 2$  for all  $x \in [0, 2]$ ?

3. Let  $f$  be twice differentiable on  $[0, 2]$ . Suppose that  $f(0) = 0, f(1) = 2$  and  $f(2) = 4$ . Show that there is  $x_0 \in (0, 2)$  such that  $f''(x_0) = 0$ .

4. Let  $a > 0$  and  $f : [-a, a] \rightarrow \mathbb{R}$  be differentiable. Suppose that  $f'(x) \leq 1$  for all  $x \in (-a, a)$ . If  $f(a) = a$  and  $f(-a) = -a$ , then show that  $f(x) = x$  for every  $x \in (-a, a)$ .

5. Let  $f : [0, 1] \rightarrow \mathbb{R}$  be twice differentiable. Suppose that the line segment joining the points  $(0, f(0))$  and  $(1, f(1))$  intersect the graph of  $f$  at a point  $(a, f(a))$  where  $0 < a < 1$ . Show that there exists  $x_0 \in [0, 1]$  such that  $f''(x_0) = 0$ .

6. Let  $f : [0, 1] \rightarrow \mathbb{R}$  be continuous. Suppose that  $f$  is differentiable on  $(0, 1)$  and  $\lim_{x \rightarrow 0} f'(x) = \alpha$  for some  $\alpha \in \mathbb{R}$ . Show that  $f'(0)$  exists and  $f'(0) = \alpha$ .

7. Let  $f : [0, 1] \rightarrow \mathbb{R}$  be differentiable and  $f(0) = 0$ . Suppose that  $|f'(x)| \leq |f(x)|$  for all  $x \in [0, 1]$ . Show that  $f(x) = 0$  for all  $x \in [0, 1]$ .

8. Let  $f : [0, \infty) \rightarrow \mathbb{R}$  be continuous and  $f(0) = 0$ . Suppose that  $f'(x)$  exists for all  $x \in (0, \infty)$  and  $f'$  is increasing on  $(0, \infty)$ . Show that the function  $g(x) = \frac{f(x)}{x}$  is increasing on  $(0, \infty)$ .

9. Establish the following inequalities.

(a) For  $\alpha > 1$ ,  $(1+x)^\alpha \geq 1 + \alpha x$  for all  $x > -1$ .

(b) For  $x > 0$ ,  $e \log x \leq x$ .

10. Let  $a > 0$  and  $f : [a, b] \rightarrow \mathbb{R}$  be differentiable. Show that there exists  $c \in (a, b)$  such that  $\frac{bf(a)-af(b)}{b-a} = f(c) - cf'(c)$ .

11. Let  $f : [a, b] \rightarrow \mathbb{R}$  be differentiable and  $a \geq 0$ . Using the Cauchy mean value theorem, show that there exist  $c_1, c_2 \in (a, b)$  such that  $\frac{f'(c_1)}{a+b} = \frac{f'(c_2)}{2c_2}$ .

12. Evaluate the following limits using L'Hospital's Rule.

(a)  $\lim_{x \rightarrow 0^+} \left( \frac{1}{\sin x} - \frac{1}{x} \right)$ .

(b)  $\lim_{x \rightarrow \infty} (e^x + x)^{1/x}$ .

(c)  $\lim_{x \rightarrow \infty} (\log x - x)$ .

(d)  $\lim_{x \rightarrow 0^+} (\cos x)^{1/x}$ .

(e)  $\lim_{x \rightarrow 0^+} (\sin x)^{\sqrt{x}}$ .

13. Let  $f : (0, \infty) \rightarrow [1, \infty)$  be differentiable. Suppose that  $\lim_{x \rightarrow \infty} (f(x) + f'(x)) = \ell$  for some  $\ell \in \mathbb{R}$ . Using L'Hospital's rule, show that  $\lim_{x \rightarrow \infty} f(x) = \ell$ .

14. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be such that  $f''(c)$  exists at some  $c \in \mathbb{R}$ . Using L'Hospital's Rule, show that

$$\lim_{h \rightarrow 0} \frac{f(c+h) - 2f(c) + f(c-h)}{h^2} = f''(c).$$

Give an example of a function  $f$  and a point  $c$  such that the above limit exists but  $f$  is not twice differentiable at  $c$ .

15. (\*) Let  $f : [a, b] \rightarrow \mathbb{R}$  be differentiable. If  $f'(x) \neq 0$  for all  $x \in [a, b]$ , then show that either  $f'(x) > 0$  for all  $x \in [a, b]$  or  $f'(x) < 0$  for all  $x \in [a, b]$ .

16. (\*) Let  $f : [a, b] \rightarrow \mathbb{R}$  be such that  $f'(x) \neq 0$  for all  $x \in [a, b]$  and  $J = \{f(x) : x \in [a, b]\}$ . Show that  $f^{-1} : J \rightarrow [a, b]$  is differentiable and  $(f^{-1})'(f(x)) = \frac{1}{f'(x)}$  for all  $x \in [a, b]$ .

Practice Problems 8: Hints/Solutions

1. (a) By the MVT, there exists  $c \in (n, n+1)$  such that  $\sqrt{n+1} - \sqrt{n} = \frac{1}{2\sqrt{c}}$ .  
 (b) By the MVT, there exists  $c \in (1, x)$  such that  $\log x - \ln 1 = \frac{1}{c}(x-1)$ .  
 (c) By the MVT (see Application 7.3),  $e^x \geq 1+x$  for all  $x \in \mathbb{R}$ . That is,  $e^{x-1} \geq 1+(x-1)$ .
2. If so, then by the MVT there exists  $c \in (0, 2)$  such that  $5 = f(2) - f(0) = 2f'(c)$ .
3. By the MVT there exist  $x_1 \in (0, 1)$  and  $x_2 \in (1, 2)$  such that  $f'(x_1) = f(1) - f(0) = 2$  and  $f'(x_2) = f(2) - f(1) = 2$ . Apply Rolle's theorem for  $f'$  on  $[x_1, x_2]$
4. Let  $g(x) = f(x) - x$  for all  $x \in [-a, a]$ . Note that  $g'(x) \leq 0$  on  $(-a, a)$ . Therefore,  $g$  is decreasing. Since  $g(a) = g(-a) = 0$ , we have  $g = 0$ .
5. Using the MVT on  $[0, a]$  and  $[a, 1]$ , obtain  $b \in (0, a)$  and  $c \in (a, 1)$  such that  $\frac{f(a)-f(0)}{a-0} = f'(b)$  and  $\frac{f(1)-f(a)}{1-a} = f'(c)$ . Note that  $f'(b) = f'(c)$  because they are slopes of the same chord. By Rolle's theorem there exists  $x_0 \in (b, c)$  such that  $f''(x_0) = 0$ .
6. Let  $x \in (0, 1]$ . By the MVT, there exists  $c_x \in (0, x)$  such that  $\frac{f(x)-f(0)}{x} = f'(c_x)$ . Now  $f'(0) = \lim_{x \rightarrow 0} \frac{f(x)-f(0)}{x} = \lim_{x \rightarrow 0} f'(c_x) = \lim_{c_x \rightarrow 0} f'(c_x) = \alpha$ .
7. For  $x \in (0, 1)$ , by the MVT, there exists  $x_1$  such that  $0 < x_1 < x$  and  $f(x) = f'(x_1)x$ . This implies that  $|f(x)| \leq x|f'(x_1)|$ . Similarly there exists  $x_2$  such that  $0 < x_2 < x_1$  and  $|f(x_1)| \leq x_1|f'(x_2)|$ . Therefore  $|f(x)| \leq x^2|f'(x_2)|$ . Find a sequence  $(x_n)$  in  $(0, 1)$  such that  $|f(x)| \leq x^n|f'(x_n)|$ . Since  $f$  is bounded on  $[0, 1]$ ,  $x^n|f'(x_n)| \rightarrow 0$ . Hence  $f(x) = 0$ .
8. Note that  $g'(x) = \frac{xf'(x)-f(x)}{x^2} = \frac{f'(x)-\frac{f(x)}{x}}{x}$ . Observe that, by the MVT,  $\frac{f(x)}{x} = f'(c_x)$  for some  $c_x \in (0, x)$ . Since  $f'$  is increasing,  $g'(x) \geq 0$ . Hence  $g$  is increasing.
9. (a) Let  $\alpha > 1$  and  $f(x) = (1+x)^\alpha - (1+\alpha x)$  on  $(-1, \infty)$ . Then  $f'(x) \leq 0$  on  $(-1, 0]$  and  $f'(x) \geq 0$  on  $[0, \infty)$ . Hence  $f(x) \geq f(0) = 0$  on  $(-1, 0]$  and  $f(x) \geq f(0) = 0$  on  $[0, \infty)$ . Therefore  $f(x) \geq 0$  on  $(-1, \infty)$ .  
 (b) Define  $f(x) = x - e \log x$  on  $(0, \infty)$ . Then  $f'(x) = \frac{x-e}{x}$ . Therefore  $f'(x) > 0$  on  $(e, \infty)$  and  $f'(x) < 0$  on  $(0, e)$ . Hence  $f(x) > f(e)$  for all  $x \in (0, \infty)$  and  $x \neq e$ .
10. Observe that  $\frac{bf(a)-af(b)}{b-a} = \frac{\frac{f(b)}{\frac{1}{b}} - \frac{f(a)}{\frac{1}{a}}}{\frac{1}{b} - \frac{1}{a}}$ . Apply the CMVT to  $\frac{f(x)}{x}$  and  $\frac{1}{x}$ .
11. Apply the CMVT to  $f(x)$  and  $g_1(x) = x$ . Again apply to  $f(x)$  and  $g_2(x) = x^2$ .
12. (a) We have  $\lim_{x \rightarrow 0^+} \left( \frac{1}{\sin x} - \frac{1}{x} \right) = \lim_{x \rightarrow 0^+} \frac{x - \sin x}{x \sin x} = \lim_{x \rightarrow 0^+} \frac{1 - \cos x}{\sin x + x \cos x} = \lim_{x \rightarrow 0^+} \frac{\sin x}{2 \cos x - x \sin x} = 0$ .  
 (b) Note that  $\log(\lim_{x \rightarrow \infty} (e^x + x)^{1/x}) = \lim_{x \rightarrow \infty} \log(e^x + x)^{1/x} = \lim_{x \rightarrow \infty} \frac{\log(e^x + x)}{x} = 1$ . Thus  $\lim_{x \rightarrow \infty} (e^x + x)^{1/x} = e$ .  
 (c) Observe that  $\log x - x = \log(xe^{-x})$  and  $\lim_{x \rightarrow \infty} \frac{x}{e^x} = 0$ . Thus  $\lim_{x \rightarrow \infty} (\log x - x) = -\infty$ .  
 (d) Since  $\log(\lim_{x \rightarrow 0^+} (\cos x)^{\frac{1}{x}}) = \lim_{x \rightarrow 0^+} \log((\cos x)^{\frac{1}{x}}) = \lim_{x \rightarrow 0^+} \frac{\log(\cos x)}{x} = \lim_{x \rightarrow 0^+} (-\tan x) = 0$ ,  $\lim_{x \rightarrow 0^+} (\cos x)^{\frac{1}{x}} = e^0 = 1$ .  
 (e) Since  $\log(\lim_{x \rightarrow 0^+} (\sin x)^{\sqrt{x}}) = \lim_{x \rightarrow 0^+} \frac{\log(\sin x)}{1/\sqrt{x}} = -2 \lim_{x \rightarrow 0^+} \frac{x^{\frac{3}{2}} \cos x}{\sin x}$  and  $\lim_{x \rightarrow 0^+} \frac{x^{\frac{3}{2}} \cos x}{\sin x} = \left( \lim_{x \rightarrow 0^+} \sqrt{x} \cos x \right) \left( \lim_{x \rightarrow 0^+} \frac{x}{\sin x} \right) = 0$ ,  $\lim_{x \rightarrow 0^+} (\sin x)^{\sqrt{x}} = e^0 = 1$ .

13. Observe that  $f(x) = \frac{e^x f(x)}{e^x}$ . Apply L'Hospital's Rule.
14. Since  $f''(c)$  exists, there exists a  $\delta > 0$  such that  $f'(x)$  exists on  $(c - \delta, c + \delta)$ . Therefore, by L'Hospital's Rule, the given limit is equal to  $\lim_{h \rightarrow 0} \frac{f'(c+h) - f'(c-h)}{2h}$  if it exists. But  $\lim_{h \rightarrow 0} \frac{f'(c+h) - f'(c-h)}{2h} = \frac{1}{2} \left[ \lim_{h \rightarrow 0} \frac{f'(c+h) - f'(c)}{h} + \lim_{h \rightarrow 0} \frac{f'(c-h) - f'(c)}{-h} \right] = \frac{1}{2} [f''(c) + f''(c)]$ .
- Let  $f(x) = 1$  on  $(0, \infty)$ ,  $f(0) = 0$  and  $f(x) = -1$  on  $(-\infty, 0)$ . Then  $f$  is not continuous at 0 hence  $f''(0)$  does not exist. It can be easily verified that the limit given in the question exists.
15. Since  $f$  is one-one (see Application 7.2), it is either strictly increasing or strictly decreasing (see Problem 20 of PP 6). Hence either  $f'(x) \geq 0$  for all  $x \in [a, b]$  or  $f'(x) \leq 0$  for all  $x \in [a, b]$ . This problem can also be solved using Problem 18 in PP 7.
16. First note that  $f$  is one-one as  $f'(x) \neq 0$  for all  $x \in [a, b]$  (See Application 7.2). Let  $y_0 \in J$  and  $y_0 = f(x_0)$  for some  $x_0 \in [a, b]$ . Let  $(y_n)$  be any arbitrary sequence in  $J$  such that  $y_n \neq y_0$  for all  $n$ ,  $y_n \rightarrow y_0$  and  $y_n = f(x_n)$  for some  $x_n \in [a, b]$ . Since  $f^{-1}$  is continuous (see Problem 16 in PP5) and  $f^{-1}$  is also one-one, we have  $x_n \rightarrow x_0$  and  $x_n \neq x_0$  for all  $n$ . Now

$$\lim_{n \rightarrow \infty} \frac{f^{-1}(y_n) - f^{-1}(y_0)}{y_n - y_0} = \lim_{n \rightarrow \infty} \frac{x_n - x_0}{f(x_n) - f(x_0)} = \lim_{n \rightarrow \infty} \frac{1}{\frac{f(x_n) - f(x_0)}{x_n - x_0}} = \frac{1}{f'(x_0)}.$$