

Name:

Roll No:

Section:

Quiz-II, MTH-102A: LINEAR ALGEBRA AND ODE

Date: 5th April, 2019

Time: 6.30 PM-7.00 PM

Max. Marks: 20

1. Write your Name, Roll No. and Section.
2. Write answers only in the space provided.
3. No additional sheets will be provided.

1. Find a suitable integrating factor and solve the equation.

$$xy' + (2 - 4x)y = \frac{e^{4x}}{x}, \quad x > 0.$$

[3]

Answer:

Solution:

For $y' + p(x)y = r(x)$, a integrating factor is $\mu(x) = e^{\int p dx}$. Here

$$p(x) = \frac{2}{x} - 4, \quad r(x) = \frac{e^{4x}}{x^2}.$$

So

$$\mu(x) = x^2 e^{-4x}.$$

[2]

Hence the solution is $y\mu(x) = \int r(x)\mu(x) = x + c$. So

$$y = c \frac{e^{4x}}{x^2} + \frac{e^{4x}}{x}.$$

[1]

2. Which pairs of functions below can be an independent set of solutions for $y'' + p(x)y' + q(x)y = 0$ with $p(x), q(x)$ being continuous on the interval as indicated below? Justify your answer in each case.

(a) $y_1 = x, \quad y_2 = \cos x, \quad x \in (0, \infty)$.

(b) $y_1 = x^3, \quad y_2 = x, \quad x \in (0, \infty)$.

(c) $y_1 = 0, \quad y_2 = e^{4x+1}, \quad x \in (-\infty, \infty)$.

[5]

Answer:

Solution: Clearly functions in (c) are linearly dependent.

[1]

For (a) the wronskian is $(-x \sin x - \cos x)$. It has zero at infinitely many points in $(0, \infty)$. So they can not be independent solutions of second order linear homogeneous ODE.

[2]

For (b) the wronskian is $(x^3 - 3x^3 = -2x^3)$. It is NEVER zero on $(0, \infty)$. So they are independent solution of second order linear homogeneous ODE.

[2]

3. Find the general solution of the following equation using variation of parameters.

$$y'' + 4y' + 4y = \frac{e^{-2x}}{x}, \quad x > 0.$$

[6]

Answer:

Solution:

• Characteristic equation of the homogeneous part : $m^2 + 4m + 4 = 0$, so $m = -2, -2$. [1]

• Two independent solution of the homogeneous part $y_1 = e^{-2x}$, $y_2 = xe^{-2x}$. [1]

• Wronskian $W(y_1, y_2) = e^{-4x}$. [1]

Particular integral $y_p = u(x)y_1 + v(x)y_2$.

• $u(x)' = -\frac{y_2 r}{W} = -1$ so $u(x) = -x$ [1]

• $v(x)' = \frac{y_1 r}{W} = 1/x$ so $v(x) = \ln(x)$. [1]

• So particular integral $y_p = -xe^{-2x} + \ln(x)xe^{-2x}$.

• General solution: $y = c_1e^{-2x} + c_2xe^{-2x} - xe^{-2x} + \ln(x)xe^{-2x}$.

or $y = c_1e^{-2x} + c_2xe^{-2x} + \ln(x)xe^{-2x}$ [1]

4. Consider the IVP

$$(x^2 - 2x)y' = 2(x - 1)y, \quad y(x_0) = y_0.$$

- (a) For which values of (x_0, y_0) , Picard's theorem implies unique solution of the the IVP?
- (b) Determine all values of (x_0, y_0) such that the IVP has no solution.
- (c) Determine all values of (x_0, y_0) such that the IVP has more than one solution. [6]

Answer:

Solution:

(a) Here $f(x, y) = 2(x - 1)y/(x^2 - 2x)$ and $\partial f/\partial y = 2(x - 1)/(x^2 - 2x)$. The existence and uniqueness theorem guarantees the existence of unique solution in the vicinity of (x_0, y_0) where f and $\partial f/\partial y$ are continuous and bounded. Thus, existence of unique solution is guaranteed at all x_0 for which $x_0(x_0 - 2) \neq 0$. Hence, unique solution exists when $x_0 \neq 0, 2$.

[2]

When $x_0 = 0$ or $x_0 = 2$, nothing can be said using the existence and uniqueness theorem. However, since the equation is separable, we can find the general solution to be $y = Cx(x - 2)$.

[2]

Using initial condition we get $y_0 = Cx_0(x_0 - 2)$. Clearly the IVP has no solution if $x_0(x_0 - 2) = 0$ and $y_0 \neq 0$.

[1]

If $x_0(x_0 - 2) = 0$ and $y_0 = 0$ then $y = \alpha x(x - 2)$ is a solution to the IVP for any real α .

[1]

Hence, in summary

- (i) No solution for $x_0 = 0$ or $x_0 = 2$ and $y_0 \neq 0$;
- (ii) Infinite number of solutions for $x_0 = 0$ or $x_0 = 2$ and $y_0 = 0$;
- (iii) Unique solution for $x_0 \neq 0, 2$.